

"Depression" in

DSM-1, DSM-2, & DSM-5

DSM-1:

000-x06 Depressive reaction

The anxiety in this reaction is allayed, and hence partially relieved, by depression and self-depreciation. The reaction is precipitated by a current situation, frequently by some loss sustained by the patient, and is often associated with a feeling of guilt for past failures or deeds. The degree of the reaction in such cases is dependent upon the intensity of the patient's ambivalent feeling toward his loss (love, possession) as well as upon the realistic circumstances of the loss.

The term is synonymous with "reactive depression" and is to be differentiated from the corresponding psychotic reaction. In this differentiation, points to be considered are (1) life history of patient, with special reference to mood swings (suggestive of psychotic reaction), to the personality structure (neurotic or cyclothymic) and to precipitating environmental factors and (2) absence of malignant symptoms (hypochondriacal preoccupation, agitation, delusions, particularly somatic, hallucinations, severe guilt feelings, intractable insomnia, suicidal ruminations, severe psychomotor retardation, profound retardation of thought, stupor).

DSM-2:

300.4 Depressive neurosis

This disorder is manifested by an excessive reaction of depression due to an internal conflict or to an identifiable event such as the loss of a love object or cherished possession. It is to be distinguished from *Involucional melancholia* (q.v.) and *Manic-depressive illness* (q.v.). *Reactive depressions* or *Depressive reactions* are to be classified here.

should receive one of those diagnoses rather than disruptive mood dysregulation disorder. Children with disruptive mood dysregulation disorder may have symptoms that also meet criteria for an anxiety disorder and can receive both diagnoses, but children whose irritability is manifest only in the context of exacerbation of an anxiety disorder should receive the relevant anxiety disorder diagnosis rather than disruptive mood dysregulation disorder. In addition, children with autism spectrum disorders frequently present with temper outbursts when, for example, their routines are disturbed. In that instance, the temper outbursts would be considered secondary to the autism spectrum disorder, and the child should not receive the diagnosis of disruptive mood dysregulation disorder.

Intermittent explosive disorder. Children with symptoms suggestive of intermittent explosive disorder present with instances of severe temper outbursts, much like children with disruptive mood dysregulation disorder. However, unlike disruptive mood dysregulation disorder, intermittent explosive disorder does not require persistent disruption in mood between outbursts. In addition, intermittent explosive disorder requires only 3 months of active symptoms, in contrast to the 12-month requirement for disruptive mood dysregulation disorder. Thus, these two diagnoses should not be made in the same child. For children with outbursts and intercurrent, persistent irritability, only the diagnosis of disruptive mood dysregulation disorder should be made.

Comorbidity

Rates of comorbidity in disruptive mood dysregulation disorder are extremely high. It is rare to find individuals whose symptoms meet criteria for disruptive mood dysregulation disorder alone. Comorbidity between disruptive mood dysregulation disorder and other DSM-defined syndromes appears higher than for many other pediatric mental illnesses; the strongest overlap is with oppositional defiant disorder. Not only is the overall rate of comorbidity high in disruptive mood dysregulation disorder, but also the range of comorbid illnesses appears particularly diverse. These children typically present to the clinic with a wide range of disruptive behavior, mood, anxiety, and even autism spectrum symptoms and diagnoses. However, children with disruptive mood dysregulation disorder should not have symptoms that meet criteria for bipolar disorder, as in that context, only the bipolar disorder diagnosis should be made. If children have symptoms that meet criteria for oppositional defiant disorder or intermittent explosive disorder and disruptive mood dysregulation disorder, only the diagnosis of disruptive mood dysregulation disorder should be assigned. Also, as noted earlier, the diagnosis of disruptive mood dysregulation disorder should not be assigned if the symptoms occur only in an anxiety-provoking context, when the routines of a child with autism spectrum disorder or obsessive-compulsive disorder are disturbed, or in the context of a major depressive episode.

Major Depressive Disorder

Diagnostic Criteria

- A. Five (or more) of the following symptoms have been present during the same 2-week period and represent a change from previous functioning; at least one of the symptoms is either (1) depressed mood or (2) loss of interest or pleasure.

DSM-5:

- Note:** Do not include symptoms that are clearly attributable to another medical condition.
1. Depressed mood most of the day, nearly every day, as indicated by either subjective report (e.g., feels sad, empty, hopeless) or observation made by others (e.g., appears tearful). (**Note:** In children and adolescents, can be irritable mood.)
 2. Markedly diminished interest or pleasure in all, or almost all, activities most of the day, nearly every day (as indicated by either subjective account or observation).

and dysregulation disorder, but children whose irritable mood dysregulation disorder should be distinguished from the disruptive mood dysregulation disorder. In that instance, the disruptive mood dysregulation disorder, and the disruptive mood dysregulation disorder.

Disruptive mood dysregulation disorder, much like children with disruptive mood dysregulation disorder, requires only 3 months of symptoms in the same child. For the diagnosis of dis-

ruptive mood dysregulation disorder and other psychiatric mental illnesses; the overall rate of the range of comorbid symptoms present to the clinic even autism spectrum disorder, as in that context, the symptoms that meet criteria for disruptive mood dysregulation disorder only in an anxiety disorder or obsessive depressive episode.

Major Depressive Disorder

During the same 2-week period, one of the symptoms

other medical condition.

Attributed by either subjective or objective (e.g., irritable mood.)

Activities most of the time (or observation).

3. Significant weight loss when not dieting or weight gain (e.g., a change of more than 5% of body weight in a month), or decrease or increase in appetite nearly every day. (Note: In children, consider failure to make expected weight gain.)
4. Insomnia or hypersomnia nearly every day.
5. Psychomotor agitation or retardation nearly every day (observable by others, not merely subjective feelings of restlessness or being slowed down).
6. Fatigue or loss of energy nearly every day.
7. Feelings of worthlessness or excessive or inappropriate guilt (which may be delusional) nearly every day (not merely self-reproach or guilt about being sick).
8. Diminished ability to think or concentrate, or indecisiveness, nearly every day (either by subjective account or as observed by others).
9. Recurrent thoughts of death (not just fear of dying), recurrent suicidal ideation without a specific plan, or a suicide attempt or a specific plan for committing suicide.

- B. The symptoms cause clinically significant distress or impairment in social, occupational, or other important areas of functioning.
- C. The episode is not attributable to the physiological effects of a substance or to another medical condition.

Note: Criteria A–C represent a major depressive episode.

Note: Responses to a significant loss (e.g., bereavement, financial ruin, losses from a natural disaster, a serious medical illness or disability) may include the feelings of intense sadness, rumination about the loss, insomnia, poor appetite, and weight loss noted in Criterion A, which may resemble a depressive episode. Although such symptoms may be understandable or considered appropriate to the loss, the presence of a major depressive episode in addition to the normal response to a significant loss should also be carefully considered. This decision inevitably requires the exercise of clinical judgment based on the individual's history and the cultural norms for the expression of distress in the context of loss.¹

- D. The occurrence of the major depressive episode is not better explained by schizoaffective disorder, schizophrenia, schizophreniform disorder, delusional disorder, or other specified and unspecified schizophrenia spectrum and other psychotic disorders.
- E. There has never been a manic episode or a hypomanic episode.

Note: This exclusion does not apply if all of the manic-like or hypomanic-like episodes are substance-induced or are attributable to the physiological effects of another medical condition.

¹In distinguishing grief from a major depressive episode (MDE), it is useful to consider that in grief the predominant affect is feelings of emptiness and loss, while in MDE it is persistent depressed mood and the inability to anticipate happiness or pleasure. The dysphoria in grief is likely to decrease in intensity over days to weeks and occurs in waves, the so-called pangs of grief. These waves tend to be associated with thoughts or reminders of the deceased. The depressed mood of MDE is more persistent and not tied to specific thoughts or preoccupations. The pain of grief may be accompanied by positive emotions and humor that are uncharacteristic of the pervasive unhappiness and misery characteristic of MDE. The thought content associated with grief generally features a preoccupation with thoughts and memories of the deceased, rather than the self-critical or pessimistic ruminations seen in MDE. In grief, self-esteem is generally preserved, whereas in MDE feelings of worthlessness and self-loathing are common. If self-derogatory ideation is present in grief, it typically involves perceived failings vis-à-vis the deceased (e.g., not visiting frequently enough, not telling the deceased how much he or she was loved). If a bereaved individual thinks about death and dying, such thoughts are generally focused on the deceased and possibly about "joining" the deceased, whereas in MDE such thoughts are focused on ending one's own life because of feeling worthless, undeserving of life, or unable to cope with the pain of depression.

DSM-5 Continued