

## How Will We Know a Lapse When We See One? Comment on Leri and Stewart (2002)

Timothy B. Baker and John J. Curtin  
University of Wisconsin—Madison

The article by F. Leri and J. Stewart (2002) addresses the validity of animal models of relapse in a more sophisticated manner than does much prior research. These researchers have shown that drug self-administration can be influenced by the presence of drug contingent cues as well as by active self-administration versus passive infusion of “lapse” doses. This research also leads to additional questions about the external validity of animal relapse models. Current relapse models may lack validity because of the parameters of drug exposure, because abstinence is imposed on the organism, and because there is no motivational influence that counters resumption of drug self-administration. F. Leri and J. Stewart’s (2002) article encourages a more thorough assessment of the motivational context of relapse models.

We were asked to provide a peer commentary with respect to the article, “The Consequences of Different ‘Lapses’ on Relapse to Heroin Seeking in Rats,” by Francesco Leri and Jane Stewart (2002). It is a pleasure to provide commentary because this article was intriguing and thought provoking.

One reason that the work is thought provoking is that it raises important questions about the fundamental nature of relapse. As the authors note, relapse is not synonymous with any resumption of drug use after a period of abstinence. Rather, a modest resumption of use that is quickly curtailed (i.e., a “lapse”) does not really capture relapse in a way that reflects its clinical or theoretical significance. Given a fundamental distinction between a lapse and relapse, the authors rightly question whether there is something about the nature of the lapse, or lapse process, that influences the trajectory of future drug use. In particular, the authors ask whether a lapse-induced acceleration of drug self-administration depends on the organism’s engaging in the instrumental act of self-administration or whether a lapse increases subsequent drug self-administration because of drug effects per se (i.e., passive infusion exerts effects similar to those produced by self-administration). In addition, the authors systematically evaluate the impact of stimulus–contextual features of the lapse on animals’ subsequent self-administration behavior.

Leri and Stewart’s (2002) results showed interesting and rather robust effects of cue contingencies and control over drug delivery on self-administration behavior. For instance, the presence of contextual stimuli previously contingent

with drug infusions was associated with increased drug self-administration during both the lapse as well as the relapse sessions. However, the presence of these stimuli during the lapse exerted relatively little impact during the subsequent drug availability (i.e., the relapse). Instead, the self-administration of drug during the relapse appeared to be influenced by a prior contingency between lever pressing and drug effects during the lapse.

What are the potential messages of this work? One message is that drug exposure, per se, does not lead to as strong a resumption of drug self-administration as does drug infusion contingent with the execution of a previously drug-consequated operant. Hence, a maximal increase in the trajectory of drug self-administration may require not just that the organism is reminded of drug effects but is reminded instead of the response–drug nexus.

Another potentially important observation of this research is that an apparently effective prod to greater drug self-administration over time is not drug exposure or drug self-administration, per se, but rather drug availability and exposure following a period of deprivation. Thus, the authors note that, across self-administration sessions, initial responding for the drug increased steadily (relative to the prior session) after each period of drug abstinence that preceded the sessions. Moreover, this boost in self-administration was greatest after the prolonged period of drug deprivation imposed by the extinction procedure. This observation joins a growing body of evidence that suggests that researchers may have too blithely jettisoned deprivation as a vital explanatory element in their attempt to account for the motivation to use addictive drugs. Although substantial evidence indicates that addictive drugs can serve as positive reinforcers and impart incentive effects, too little attention has been directed at how negative and positive reinforcement processes may interact in affecting the future incentive value of the drug or how deprivation, per se, may inflate future incentive value. Thus, like other restorative or repleting responses, what is crucial is not that the organism experiences drug effects and the effects of deprivation but

---

Timothy B. Baker and John J. Curtin, Center for Tobacco Research and Intervention, University of Wisconsin—Madison.

This work was supported by Grant NCI 1 P50 CA84724 from the National Cancer Institute.

Correspondence concerning this article should be addressed to Timothy B. Baker, Center for Tobacco Research and Intervention, University of Wisconsin, 1930 Monroe Street, Suite 200, Madison, Wisconsin 53711. E-mail: tbb@cetri.medicine.wisc.edu



study). One typically thinks of relapse as occurring after some period in which the organism has desisted from drug seeking.

Perhaps the most significant concern with respect to animal models of relapse is that researchers have not modeled the interplay of motivational influences that occurs over the course of addictive drug use in humans, and that sets the stage for abstinence. These motivational forces must be overcome or neutralized for relapse to occur. That is, the authors never created a psychological or motivational context that produced drug abstinence.

Without the creation of countervailing motives, relapse research becomes the study of availability that is, in the Leri and Stewart (2002) study, all that animals required in order to resume significant drug self-administration was information that drug was once more available via lever pressing. Although information on drug availability can exert potent effects in human addicts (Juliano & Brandon, 1998; Zinzer et al., 1999), we wonder whether this is indeed a crucial determinant of relapse. How many smokers trying to quit do not know that nicotine is readily available from the nearest convenience store or obliging tobacco user? Indeed, why should an organism not resume the self-administration of any appetitive commodity when it yields pleasure and produces no cost? It is unlikely that the rat has any appreciation of the ultimate costs of addiction, just as William James's "broody hen" had no prophetic inkling of the results of sitting on eggs (James, 1890/1950).

In sum, one of the defining features of relapse may be the resumption of drug use despite strong reasons not to do so, for example, punishment, or anticipated punishment, for "falling off the wagon" (see Breiner, Stritzke, & Lang, 1999). Ignoring this press for abstinence could mislead researchers in their attempt to characterize the nature of relapse precipitants. Let us assume for the sake of argument that addictive drug motivation is multifactorial (e.g., it reflects the impact of positive reinforcement, negative reinforcement, and overconfidence in estimates of control). It may be, in fact, that there is a hierarchy of motives, with some producing particularly compelling motivation. Cues and contingencies that may be efficacious signals of availability may be ineffective in the presence of strong motives against drug use. In short, we are suggesting that by ignoring the psychological features of the relapse situation, we may be casting too wide a net, capturing dross along with potent relapse precipitants.

In conclusion, it may be that researchers have not paid sufficient attention to the behavioral and psychopharmacological processes that characterize clinical phenomena such as lapses and relapses. The work of Leri and Stewart (2002) is an important step toward a more careful analysis and unpacking of these phenomena.

## References

- Baker, T. B., Morse, E., & Sherman, J. E. (1987). *The motivation to use drugs: A psychobiological analysis of urges*. In C. Rivers (Ed.), *Nebraska Symposium on Motivation: Vol. 34. Alcohol use and abuse* (pp. 257-323). Lincoln: University of Nebraska Press.
- Bickel, W. K., Stitzer, M. L., Liebson, I. A., & Bigelow, G. E. (1988). Acute physical dependence in man: Effects of naloxone after brief morphine exposure. *Journal of Pharmacology and Experimental Therapeutics*, *133*, 262-270.
- Brandon, T. H., Tiffany, S. T., Obremski, K. M., & Baker, T. B. (1990). Postcessation cigarette use: The process of relapse. *Addictive Behaviors*, *15*, 105-114.
- Breiner, M. J., Stritzke, W. G. K., & Lang, A. R. (1999). Approaching avoidance: A step essential to the understanding of craving. *Alcohol Research and Health*, *23*, 197-206.
- Hall, W. G., Arnold, H. M., & Myers, K. P. (2000). The acquisition of an appetite. *Psychological Science*, *11*, 101-105.
- Heischman, S. J., Stitzer, M. L., Bigelow, G. E., & Liebson, I. A. (1989). Acute opioid physical dependence in postaddict humans: Naloxone dose effects after brief morphine exposure. *The Journal of Pharmacology and Experimental Therapeutics*, *248*, 127-134.
- James, W. (1950). *The principles of psychology*. New York: Dover. (Original work published 1890)
- Juliano, L. M., & Brandon, T. H. (1998). Reactivity to instructed smoking availability and environmental cues: Evidence with urge and reaction time. *Experimental and Clinical Psychopharmacology*, *6*, 45-53.
- Leri, F., & Stewart, J. (2002). The consequences of different "lapses" on relapse to heroin seeking in rats. *Experimental and Clinical Psychopharmacology*, *10*, 000-000.
- Lyvers, M. (1998). Drug addiction as a physical disease: The role of physical dependence and other chronic drug-induced neurophysiological changes in compulsive drug self-administration. *Experimental and Clinical Psychopharmacology*, *6*, 107-125.
- Martin, W. R., & Eades, C. G. (1961). Demonstration of tolerance and physical dependence in the dog following a short-term infusion of morphine. *Journal of Pharmacology and Experimental Therapeutics*, *133*, 262-270.
- Pomerleau, C. S., & Pomerleau, O. F. (1992). Euphoriant effects of nicotine in smokers. *Psychopharmacology*, *108*, 460-465.
- Robinson, T. E., & Berridge, K. C. (1993). The neural basis of drug craving: An incentive-sensitization theory of addiction. *Brain Research Reviews*, *18*, 247-291.
- Stewart, J., de Wit, H., & Eikelboom, R. (1984). Role of unconditioned and conditioned drug effects in the self-administration of opiates and stimulants. *Psychological Review*, *91*, 251-268.
- Ying, F. (1998). *Smoking motivation, nicotine withdrawal and the startle probe response*. Unpublished doctoral dissertation, University of Wisconsin—Madison.
- Zinzer, M. C., Fiore, M. C., Davidson, R. J., & Baker, T. B. (1999). Manipulating smoking motivation: Impact on an electrophysiological index of approach motivation. *Journal of Abnormal Psychology*, *108*, 240-254.

Received January 16, 2002

Accepted January 23, 2002 ■